



A complete course of treatment for trichomoniasis (trich) with **just one dose**¹



Effective treatment for trich is available in a **single dose**—not multiple days of pills^{1,2}

One day.
One dose.
That's it!



About Trich³

- Like all sexually transmitted infections (STIs), trich is **passed from partner to partner** during sex
- About **70%** of people infected with trich have **no symptoms**
- **Left untreated**, trich in women is associated with increased risk of:

Getting other STIs, including herpes, chlamydia, gonorrhea, and HIV, and transmitting STIs to your partner ^{3,4}	Developing pelvic inflammatory disease (PID), which can threaten your fertility ^{5,6}	Developing cervical cancer, which can be life-threatening ⁷	Complications during pregnancy, including pre-term birth ³
--	---	---	--

You should know

- Since trich can last for **months or even years** without treatment, a trich diagnosis does not necessarily mean you were recently infected³

Treating Trich

- Since trich is an STI, **you and your partner(s)** should be treated with antibiotics **at the same time** to prevent reinfection³
 - In some states, your healthcare provider can write a prescription for your partner(s) without an examination^{8,9}
- Don't have sexual intercourse with your partner(s) until you are all treated for trich³

With SOLOSEC, you and your partner(s) can complete treatment for trich with **just one oral dose**

What is SOLOSEC?

SOLOSEC® (secnidazole) 2g oral granules is a prescription medicine used to treat bacterial vaginosis in adult women. SOLOSEC is used to treat Trichomoniasis, a common sexually transmitted infection (STI), in adults. People should avoid having sex until they and their sex partners are treated with the same dose and at the same time to prevent reinfection.

How should I take SOLOSEC?

SOLOSEC is a single-dose therapy for oral use. Use SOLOSEC by sprinkling an entire packet of SOLOSEC onto applesauce, yogurt, or pudding. The entire dose should be taken at once, and finished within 30 minutes. Avoid chewing or crunching the granules. SOLOSEC should not be taken by dissolving the granules in any liquid. Avoid drinking alcohol during treatment with SOLOSEC and for 2 days (48 hours) after you take SOLOSEC.

See additional Important Safety Information continued on reverse side.

Questions to Ask Your Healthcare Provider Before You Leave

- ✓ What should I tell my partner(s) about my trich diagnosis?
- ✓ What kind of treatment do partner(s) need, and when should they take it?
- ✓ Are you able to provide a prescription for my partner(s) directly?
- ✓ If not, can you supply me with information for my partner(s), or a referral for treatment?
- ✓ What can I do to keep from being reinfected with trich?

For answers to common questions about trich and information about treatment with SOLOSEC, visit [solosec.com](https://www.solosec.com)

One day.
One dose.
That's it!



To take SOLOSEC, mix the entire packet with your choice of yogurt, pudding, or applesauce and eat the mixture within 30 minutes without chewing or crunching the granules.

Treatment with SOLOSEC is easier to complete with one dose in one day, compared to other medicines that may require as many as 14 pills over 7 days^{1,2}



SOLOSEC 2 g
1 Dose, ONCE



Another common antibiotic for trich
14 Doses, 2x per day for 7 days

Important Safety Information

- You should not use SOLOSEC if you've had an allergic reaction to secnidazole, other ingredients of the formulation, or other nitroimidazole derivatives.
- Before taking SOLOSEC, tell your healthcare provider about all of your medical conditions, including if you
 - Are pregnant or plan to become pregnant.
 - Are breastfeeding or plan to breastfeed. Breastfeeding is not recommended. You should not breastfeed for 96 hours (4 days) after taking SOLOSEC.
- Vaginal yeast infections may occur with SOLOSEC and require an antifungal treatment.
- The most common side effects of SOLOSEC include headache, nausea, vomiting, diarrhea, abdominal pain, and vaginal itching and a bad, bitter or metallic taste in your mouth (dysgeusia).
- Take SOLOSEC exactly as your healthcare provider tells you to. Do not take SOLOSEC more often than it is prescribed.

Call your doctor for medical advice on side effects. You are encouraged to report suspected side effects of prescription drugs to FDA at www.fda.gov/MedWatch also at 1-800-FDA-1088 or contact Lupin Pharmaceuticals, Inc. at 1-844-SOLOSEC (1-844-765-6732).

Please see full prescribing information for SOLOSEC, available at [solosec.com](https://www.solosec.com).

1. SOLOSEC [prescribing information]. Baltimore, MD: Lupin Pharmaceuticals, Inc.; 2021. 2. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin: Clinical Management Guidelines for Obstetricians-Gynecologists, Number 215. *Obstet Gynecol* 2020;135(1):e1-e17. 3. CDC Fact Sheet – Trichomoniasis. Reviewed 19 Jan 2021. Accessed 14 July 2021. 4. Kissinger P. Trichomonas vaginalis: a review of epidemiologic, clinical and treatment issues. *BMC Infect Dis* 2015;15(307):1-8. 5. Wiringa AE, Ness RB, Darville T et al. Trichomonas vaginalis, endometritis and sequelae among women with clinically suspected pelvic inflammatory disease. *Sex Transm Infect* 2020;96:436-438. 6. Workowski KA, Bolan GA; CDC. Sexually Transmitted Diseases Treatment Guidelines, 2015 [published correction appears in *MMWR Recomm Rep*. 2015;64(33):924]. *MMWR Recomm Rep* 2015;64(RR-03):1-137. 7. Yang S, Zhao W, Wang H, et al. Trichomonas vaginalis infection-associated risk of cervical cancer: A meta-analysis. *Euro J Obstet Gynecol* 2018;228:166-173. 8. Centers for Disease Control and Prevention, Division of STD Prevention. Legal Status of Expedited Partner Therapy (EPT). Updated April 2021. 9. Lupin Pharmaceuticals, Inc. Expedited Partner Therapy Legal Review. Data on File. 2021.